



Properties of Modified Double Laplace Transforms and Special Functions

V. K. Borawake^{*1} and A. P. Hiwarekar²

¹Department of Mathematics, Sinhgad Technical Education Society's Smt. Kashibai Navale College of Engineering (affiliated to Savitribai Phule Pune University), Vadgaon (Budruk), Pune 411041, Maharashtra, India

²Department of Mathematics, Vidya Pratishthan's Kamalnayan Bajaj Institute of Engineering and Technology (affiliated to Savitribai Phule Pune University), Baramati 413133, Pune, Maharashtra, India

*Corresponding author: vijayborawake@sinhgad.edu

Received: March 14, 2024

Accepted: June 6, 2024

Abstract. This paper deals with new results on modified double Laplace transforms and special functions. Starting with basic definitions and results, we have obtained a modified double Laplace transform of unit step functions, and periodic functions and developed new theorems. Finally, we illustrate our results with examples.

Keywords. Laplace transform, Multiple integral transforms, Integral transform of special functions, Double Laplace transform, Convolution

Mathematics Subject Classification (2020). 44A10, 44A20, 44A30, 44A35

Copyright © 2024 V. K. Borawake and A. P. Hiwarekar. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Laplace transform is one of the most useful transform techniques to solve various problems in mathematics and other areas. It has applications in Science, Engineering, and Technology (Nyeo *et al.* [23], Nozhak and Paskar [22]), Finance (Kim *et al.* [20], Kim and Kim [19], Daci and Tola [7]), Population growth (Daci and Tola [8]). The Laplace transform approach is a practical method for engineers, Debnath and Bhatta [9], and is used to solve various differential equations, Borawake and Hiwarekar [3], Ali *et al.* [1]. It has a wide range of applications in many fields including mathematics, physics, statistics, Poularikas and Seely [24].

The Laplace transform of a function of one variable and its applications are found in the literature, Kokulan and Lai [21], Chiu and Li [5], Chung *et al.* [6], Schiff [27], Debnath and Bhatta [9], Eltayeb and Kiliçman [15], and Hiwarekar [18]. However, such few results are available on functions of two variables and there is a need to extend this theory. The function of two or more variables and its double Laplace transform plays an important role in solving many problems, Debnath [10], Dhunde and Waghmare [11], Dhunde *et al.* [12] to solve space time-fractional equation with initial and boundary conditions, further, its applications are found in Eltayeb and Kiliçman [13–15], Tsaur and Wang [28], Hiwarekar [16–18], Riekestyn's [25], and Viaggiu [29]. Generalization of the double Laplace transforms will play an important role in developing new theory and its applications. We have extended the theory of the double Laplace transform developed by Debnath [10], and Borawake and Hiwarekar [3, 4] by obtaining new results.

2. Notations, Definitions and Basic Results

Definition 2.1 (Modified Laplace Transform). The modified Laplace transform of piece wise continuous and exponential order function $u(x)$ is

$$L_{1,a}[u(x)] = \bar{u}(p) = \int_0^{\infty} a^{-px} u(x) dx \quad (\operatorname{Re}(p) > 0, a \in (0, \infty) \setminus 1), \quad (2.1)$$

and $L_{1,a}[u(y)] = \bar{u}(q)$, provided that the integral exists (Saif *et al.* [26]), and the corresponding inverse transform is

$$L_{1,a}^{-1}[\bar{u}(p)] = \frac{1}{2\pi i} \int_{m-i\infty}^{m+i\infty} a^{px} U(p, a) dx \quad (m \geq 0).$$

Definition 2.2 (Double Laplace Transform; Debnath [10]). The double Laplace transform is given by

$$L_2[u(x, y)] = L[L[u(x, y); x \rightarrow p]; y \rightarrow q] = \int_0^{\infty} \int_0^{\infty} e^{-(px+qy)} u(x, y) dx dy \quad (2.2)$$

and the corresponding inverse transform is $L_2^{-1}[\bar{\bar{u}}(p, q)] = u(x, y)$ is defined by

$$u(x, y) = L_2^{-1}[\bar{\bar{u}}(p, q)] = \frac{1}{2\pi i} \int_{m-i\infty}^{m+i\infty} e^{px} dp \frac{1}{2\pi i} \int_{n-i\infty}^{n+i\infty} e^{qy} \bar{\bar{u}}(p, q) dq \quad (m, n \geq 0).$$

Definition 2.3 (Modified Double Laplace Transform; Borawake and Hiwarekar [3, 4]). The modified double Laplace transform of a function $u(x, y)$ is defined by

$$\bar{\bar{u}}(p, q) = L_{2,a}[u(x, y)] = L_a[L_a[u(x, y); x \rightarrow p]; y \rightarrow q] = L_{2,a}[\bar{u}(p, y); y \rightarrow q]$$

and

$$\bar{\bar{u}}(p, q) = \int_0^{\infty} \int_0^{\infty} a^{-(px+qy)} u(x, y) dx dy \quad (a > 0). \quad (2.3)$$

The modified double Laplace transform of $u(x, y)$ exists for all p and q , where $\operatorname{Re}(p) > c$ and $\operatorname{Re}(q) > d$ and $u(x, y)$ is a piece-wise continuous and of exponential order defined in finite intervals $(X, 0)$ and $(0, Y)$.

The corresponding inverse transform is

$$L_{2,a}^{-1}[\bar{\bar{u}}(p, q)] = \frac{1}{2\pi i} \int_{m-i\infty}^{m+i\infty} a^{px} dp \frac{1}{2\pi i} \int_{n-i\infty}^{n+i\infty} a^{qy} \bar{\bar{u}}(p, q) dq \quad (m, n \geq 0, a > 0).$$

Definition 2.4 (Heaviside Unit Step Function). The $H(x, y)$ is Heaviside unit step function given by

$$H(x - h, y - k) = \begin{cases} 1, & \text{if } x \geq h, y \geq k, \\ 0, & \text{if } x < h, y < k. \end{cases} \quad (2.4)$$

Definition 2.5 (Periodic Function; Amberkhane *et al.* [2]). The function $\mathbb{U}(x, y)$ is a periodic function of periods T and S given by

$$\mathbb{U}(x + T, y + S) = \mathbb{U}(x, y), \quad \text{for all } x \text{ and } y, \quad (2.5)$$

where T and S are non-zero constants and independent of x and y , respectively.

In this work, we used all terms, definitions, and standard results developed in [3, 4]. We also used the following results.

Theorem 2.1 (Shifting Property).

$$L_{2,a}[e^{-\alpha x - \beta y} u(x, y)] = L_{2,e}[p \log a + \alpha, q \log a + \beta]. \quad (2.6)$$

Theorem 2.2 (Change of Scale Property). If $L_{2,a}[u(x, y)] = \bar{u}(p, q)$, then

$$L_{2,a}[u(\alpha x, \beta y)] = \frac{1}{\alpha \beta} L_{2,a} \left[u \left(\frac{x}{\alpha}, \frac{y}{\beta} \right) \right]. \quad (2.7)$$

Here we developed the following properties of modified double Laplace transform.

3. Properties of Modified Double Laplace Transforms

In continuation with results in [3], [4] and [10], in this paper, we developed some new results on the modified double Laplace transform which are included in this section.

We consider $u(x, y)$ to be an exponentially ordered and piece-wise continuous function.

Theorem 3.1.

$$(i) \quad L_{2,a}[u(x)] = \frac{1}{q \log a} [\bar{u}(p)], \quad (3.1)$$

$$(ii) \quad L_{2,a}[u(y)] = \frac{1}{p \log a} [\bar{u}(q)]. \quad (3.2)$$

Proof. By Definition 2.3, we have

$$\begin{aligned} L_{2,a}[u(x)] &= \int_0^\infty \int_0^\infty a^{-(px+qy)} u(x) dx dy \\ &= \left[\int_0^\infty a^{-qy} dy \right] \left[\int_0^\infty a^{-px} u(x) dx \right] \\ &= \frac{1}{q \log a} [\bar{u}(p)]. \end{aligned}$$

Similarly, we have proof of (ii) part of Theorem 3.1. □

Theorem 3.2.

$$L_{2,a}[u(x + y)] = \frac{1}{(p - q) \log a} [\bar{u}(q) - \bar{u}(p)]. \quad (3.3)$$

Proof. By Definition 2.3, we have

$$\begin{aligned}
 L_{2,a}[u(x+y)] &= \int_0^\infty \int_0^\infty a^{-(px+qy)} u(x+y) dx dy \\
 &\quad (\text{Put } x+y=t, dy=dt, \text{ when } y=0, t=x \text{ and } y=\infty, t=\infty) \\
 &= \int_0^\infty a^{-(p-q)x} \left[\int_x^\infty a^{-qt} u(t) dt \right] dx.
 \end{aligned}$$

Using the change of order of integration, we have

$$\begin{aligned}
 &= \int_0^\infty \left[\int_0^t a^{-(p-q)x} dx \right] a^{-qt} u(t) dt \\
 &= \frac{1}{(p-q)\log a} \int_0^\infty [1 - a^{-(p-q)t}] a^{-qt} u(t) dt \\
 &= \frac{1}{(p-q)\log a} [\bar{u}(q) - \bar{u}(p)]. \quad \square
 \end{aligned}$$

Theorem 3.3.

$$L_{2,a}[u(x-y)] = \frac{1}{(p+q)\log a} [\bar{u}(p) + \bar{u}(q)], \text{ when } u \text{ is even.} \tag{3.4}$$

$$= \frac{1}{(p+q)\log a} [\bar{u}(p) - \bar{u}(q)], \text{ when } u \text{ is odd.} \tag{3.5}$$

Proof. By Definition 2.3, we have

$$\begin{aligned}
 L_{2,a}[u(x-y)] &= \int_0^\infty \int_0^\infty a^{-(px+qy)} u(x-y) dx dy \\
 &\quad (\text{Put } x-y=t, dy=-dt, \text{ when } y=0, t=x \text{ and } y=\infty, t=-\infty) \\
 &= \int_0^\infty a^{-px} \left[\int_{-\infty}^x a^{-q(x-t)} u(t) dt \right] dx.
 \end{aligned}$$

Using the change of order of integration, we have

$$\begin{aligned}
 &= \int_0^\infty a^{-(p+q)x} \left[\int_{-\infty}^0 a^{qt} u(t) dt \right] dx + \int_t^\infty a^{-(p+q)x} \left[\int_0^{-\infty} a^{qt} u(t) dt \right] dx \\
 &= \frac{1}{(p+q)\log a} \left[\int_{-\infty}^0 a^{qt} u(t) dt + \int_0^\infty a^{-pt} u(t) dt \right]
 \end{aligned}$$

Put $t = -\theta, dt = -d\theta$, in the first integral

$$\begin{aligned}
 &= \frac{1}{(p+q)\log a} \left[\int_\infty^0 a^{q(-\theta)} u(-\theta)(-d\theta) + \int_0^\infty a^{-pt} u(t) dt \right] \\
 &= \frac{1}{(p+q)\log a} \left[\int_0^\infty a^{q(-\theta)} u(-\theta) d\theta + \int_0^\infty a^{-pt} u(t) dt \right];
 \end{aligned}$$

$$L_{2,a}[u(x-y)] = \frac{1}{(p+q)\log a} [\bar{u}(p) + \bar{u}(q)], \text{ when } u \text{ is even.}$$

$$= \frac{1}{(p+q)\log a} [\bar{u}(p) - \bar{u}(q)], \text{ when } u \text{ is odd.} \quad \square$$

4. Modified Double Laplace Transform of Special Functions

Here we obtained the modified double Laplace transform of special functions included in the following.

Theorem 4.1 (Unit Step Function).

$$L_{2,a}[H(x - \alpha, y - \beta)] = \frac{1}{pq(\log a)^2} a^{-(p\alpha + q\beta)}, \tag{4.1}$$

where $H(x, y)$ is defined by equation (2.4) and $pq(\log a)^2 > 0$.

Proof. By Definition 2.3, we have

$$L_{2,a}[H(x - \alpha, y - \beta)] = \int_{\alpha}^{\infty} \int_{\beta}^{\infty} a^{-(px + qy)} H(x - \alpha, y - \beta) dx dy.$$

By Definition 2.4, we have

$$\begin{aligned} &= \int_{\alpha}^{\infty} \int_{\beta}^{\infty} a^{-(px + qy)} 1 dx dy \\ &= \left(\int_{\alpha}^{\infty} a^{-px} dx \right) \left(\int_{\beta}^{\infty} a^{-qy} dy \right) \\ &= \frac{1}{pq(\log a)^2} a^{-(p\alpha + q\beta)}. \end{aligned} \tag{□}$$

Theorem 4.2.

$$L_{2,a}[u(x)H(x - y)] = \frac{1}{q \log a} [\bar{u}(p) - \bar{u}(p + q)]. \tag{4.2}$$

Proof. By Definition 2.3, we have

$$\begin{aligned} L_{2,a}[u(x)H(x - y)] &= \int_0^{\infty} \int_0^{\infty} a^{-(px + qy)} u(x)H(x - y) dx dy \\ &= \int_0^{\infty} a^{-qy} \left[\int_0^{\infty} a^{-px} u(x)H(x - y) dx \right] dy \\ &= \int_0^{\infty} a^{-qy} \left[\int_y^{\infty} a^{-px} u(x) dx \right] dy. \end{aligned}$$

Using the change of order of integration, we have

$$\begin{aligned} &= \int_0^{\infty} a^{-px} u(x) \left[\int_0^x a^{-qy} dy \right] dx \\ &= \frac{1}{q \log a} \int_0^{\infty} a^{-px} u(x) [1 - a^{-qx}] dx \\ &= \frac{1}{q \log a} \left[\int_0^{\infty} a^{-px} u(x) dx - \int_0^{\infty} a^{-(p+q)x} u(x) dx \right] \\ &= \frac{1}{q \log a} [\bar{u}(p) - \bar{u}(p + q)]. \end{aligned} \tag{□}$$

Theorem 4.3.

$$L_{2,a}[u(x)H(y - x)] = \frac{1}{q \log a} [\bar{u}(p + q)]. \tag{4.3}$$

Theorem 4.4.

$$L_{2,a}[u(x)H(x+y)] = \frac{1}{q \log a} [\bar{u}(p)]. \tag{4.4}$$

Proof. By Definition 2.3, we have

$$\begin{aligned} L_{2,a}[u(x)H(x+y)] &= \int_0^\infty \int_0^\infty a^{-(px+qy)} u(x)H(x+y) dx dy \\ &= \int_0^\infty \int_0^\infty a^{-(px+qy)} u(x) dx dy \\ &= \int_0^\infty a^{-qy} dy \left[\int_0^\infty a^{-px} u(x) dx \right] \\ &= \frac{1}{q \log a} [\bar{u}(p)]. \end{aligned} \tag{4.4} \quad \square$$

Theorem 4.5.

$$L_{2,a}[H(x-y)] = \frac{1}{p(p+q)(\log a)^2}. \tag{4.5}$$

Proof. By Definition 2.3, we have

$$\begin{aligned} L_{2,a}[H(x-y)] &= \int_0^\infty \int_0^\infty a^{-(px+qy)} H(x-y) dx dy \\ &= \int_0^\infty \int_0^\infty a^{-(px+qy)} dx dy. \end{aligned}$$

Using the change of order of integration, we have

$$\begin{aligned} &= \int_0^\infty \left[\int_0^x a^{-px-xy} dy \right] dx \\ &= \frac{1}{q \log a} \int_0^\infty a^{-px} [1 - a^{-qx}] dx \\ &= \frac{1}{p(p+q)(\log a)^2}. \end{aligned} \tag{4.5} \quad \square$$

Theorem 4.6. If $\mathbb{U}(x,y)$ be a periodic function of periods T and S (Definition 2.5), and $L_{2,a}[\mathbb{U}(x,y)]$ exists, then

$$L_{2,a}[\mathbb{U}(x,y)] = \frac{1}{[1 - a^{-(pT+qS)}]} \int_0^T \int_0^S a^{-(px+qy)} \mathbb{U}(x,y) dx dy, \tag{4.6}$$

where $(1 - a^{-(pT+qS)}) > 0$.

Proof. By Definition 2.3, we have

$$\begin{aligned} L_{2,a}[\mathbb{U}(x,y)] &= \int_0^\infty \int_0^\infty a^{-(px+qy)} \mathbb{U}(x,y) dx dy \\ &= \int_0^T \int_0^S a^{-(px+qy)} \mathbb{U}(x,y) dx dy + \int_T^{2T} \int_S^{2S} a^{-(px+qy)} \mathbb{U}(x,y) dx dy \\ &\quad + \int_{2T}^{3T} \int_{2S}^{3S} a^{-(px+qy)} \mathbb{U}(x,y) dx dy + \dots \end{aligned}$$

Put $x = t + T$, $y = s + S$ in the second integral $x = t + 2T$, $y = s + 2S$ in the third integral, we have

$$L_{2,a}[\mathbb{U}(x,y)] = \int_0^T \int_0^S a^{-px-ky} \mathbb{U}(x,y) dx dy + \int_0^T \int_0^S a^{-p(t+T)-q(s+S)} \mathbb{U}(t+T,s+S) dt ds + \int_0^T \int_0^S a^{-p(t+2T)-q(s+2S)} \mathbb{U}(t+2T,s+2S) dt ds + \dots$$

Since

$$\begin{aligned} \mathbb{U}(t,s) &= \mathbb{U}(t+T,s+S) = \mathbb{U}(t+2T,s+2S) = \dots \\ &= \int_0^T \int_0^S a^{-px-ky} \mathbb{U}(x,y) dx dy + a^{-(pT+qS)} \int_0^T \int_0^S a^{-pt-qs} \mathbb{U}(t,s) dt ds \\ &\quad + a^{-2(pT+qS)} \int_0^T \int_0^S a^{-pt-qs} \mathbb{U}(t,s) dt ds + \dots \\ &= \int_0^T \int_0^S a^{-px-ky} \mathbb{U}(x,y) dx dy + a^{-(pT+qS)} \int_0^T \int_0^S a^{-px-ky} \mathbb{U}(x,y) dx dy \\ &\quad + a^{-2(pT+qS)} \int_0^T \int_0^S a^{-px-ky} \mathbb{U}(x,y) dx dy + \dots \\ &= [1 + a^{-(pT+qS)} + a^{-2(pT+qS)} + \dots] \int_0^T \int_0^S a^{-px-ky} \mathbb{U}(x,y) dx dy \\ &= \frac{1}{[1 - a^{-(pT+qS)}]} \int_0^T \int_0^S a^{-(px+ky)} \mathbb{U}(x,y) dx dy. \quad \square \end{aligned}$$

Remark 4.1. Results of Debnath [10], see equations (36), (37), (38), (39), (40), (41), (42), (43) and Theorem 3.2 are special cases of our results Theorem 3.1, Theorem 3.2, Theorem 3.3, Theorem 4.2, Theorem 4.3, Theorem 4.4, Theorem 4.5 and Theorem 4.6 respectively with $a = e$.

Remark 4.2. If we put $u(x) = 1$, in Theorem 4.5, which is a special case of Theorem 4.2.

5. Illustrative Examples

Now we illustrate our results through the following examples.

5.1: Using Theorem 3.1, equation (3.1), we have

$$L_{2,a}[\cos 2x] = \frac{p}{q(p^2 + 4)\log a}. \tag{5.1}$$

5.2: Using Theorem 3.1, equation (3.2), we have

$$L_{2,a}[\sinh(3y)] = \frac{3}{p \log a [q^2(\log a)^2 - 9]}. \tag{5.2}$$

5.3: Using Theorem 4.2, equation (4.2), we have

$$L_{2,a}[e^{2x}H(x-y)] = \frac{1}{(p \log a - 2)[(p+q)\log a - 2]}. \tag{5.3}$$

5.4: Using Theorem 4.3, equation (4.3) and Theorem 2.1, we have

$$L_{2,a}[e^{-3x}H(y-x)] = -\frac{1}{q(p+q+9)(\log a)^2}. \tag{5.4}$$

5.5: Using Theorem 4.4, equation (4.4), we have

$$L_{2,a}[x^2H(x+y)] = \frac{2}{p^3q \log a}. \quad (5.5)$$

5.6: Using Theorem 4.4, equation (4.4) and Theorem 2.2, we have

$$L_{2,a}[\sin 3xH(x+y)] = \frac{3}{(q \log a)[p^2(\log a)^2 + 9]}. \quad (5.6)$$

6. Concluding Remark

We developed new properties, theorems on modified double Laplace transform with suitable examples. There is lot of scope to extend the theory further and its corresponding applications.

Acknowledgements

V. K. Borawake is thankful to the Principal Dr. A. V. Deshpande of Sinhgad Technical Education Society's Smt. Kashibai Navale College of Engineering, Pune, and S. P. College Pune (Research Center of Mathematics), and A. P. Hiwarekar is thankful to the Principal, VPKBIET, Baramati, and to the management of Vidya Pratishthan Baramati for the entire support to this work.

Competing Interests

The authors declare that they have no competing interests.

Authors' Contributions

All the authors contributed significantly in writing this article. The authors read and approved the final manuscript.

References

- [1] A. Ali, Abdullah and A. Ahmad, The solution of Poisson partial differential equations via Double Laplace Transform Method, *Partial Differential Equations in Applied Mathematics* **4** (2021) 100058, DOI: 10.1155/2013/932578.
- [2] N. S. Amberkhane, S. P. Hatkar and K. L. Bondar, Exponential transform of some special functions, *International Journal of Mathematics Trends and Technology* **68**(6) (2022), 54 – 59, DOI: 10.14445/22315373/IJMTT-V68I6P506.
- [3] V. K. Borawake and A. P. Hiwarekar, Modified double Laplace transform of partial derivatives and its applications, *Gulf Journal of Mathematics* **16**(2) (2024), 353 – 363, DOI: 10.56947/gjom.v16i2.1892.
- [4] V. K. Borawake and A. P. Hiwarekar, New results on the modified double Laplace transform and its properties, *Journal of the Maharaja Sayajirao University of Baroda* **57**(VII) (2023), 61 – 66.
- [5] K.-S. Chiu and T. Li, Oscillatory and periodic solutions of differential equations with piecewise constant generalized mixed arguments, *Mathematische Nachrichten* **292**(10) (2019), 2153 – 2164, DOI: 10.1002/mana.201800053.
- [6] W. S. Chung, T. Kim and H. I. Know, On the q -analog of the Laplace transform, *Russian Journal of Mathematical Physics* **21** (2014), 156 – 168, DOI: 10.1134/S1061920814020034.

- [7] A. Daci and S. Tola, Application of Laplace transform in Finance, *Mathematical Modeling* **2**(4) (2018), 130 – 133, URL: <https://stumejournals.com/journals/mm/2018/4/130>.
- [8] A. Daci and S. Tola, Laplace transform, application in population growth, *International Journal of Recent Technology and Engineering* **8**(2) (2019), 954 – 957, DOI: 10.35940/ijrte.B1189.0782S419.
- [9] L. Debnath and D. Bhatta, *Integral Transform and Their Applications*, 2nd edition, Chapman and Hall/CRC, New York, 728 pages (2006), DOI: 10.1201/9781420010916.
- [10] L. Debnath, The double Laplace transforms and their properties with applications to functional, integral and partial differential equations, *International Journal of Applied and Computational Mathematics* **2** (2016), 223 – 241, DOI: 10.1007/s40819-015-0057-3.
- [11] R. R. Dhunde and G. L. Waghmare, Double Laplace transform method in mathematical physics, *International Journal of Theoretical and Mathematics Physics* **7**(1) (2017), 14 – 20, DOI: 10.5923/j.ijtmp.20170701.04.
- [12] R. R. Dhunde, N. M. Bhonde and P. R. Dhongle, Some remarks on the properties of double Laplace transforms, *International Journal of Applied Physics and Mathematics* **3**(4) (2013), 293 – 295, DOI: 10.7763/IJAPM.2013.V3.224.
- [13] H. Eltayeb and A. Kiliçman, A note on double Laplace transform and telegraphic equations, *Abstract and Applied Analysis* **2023**(1) (2013), 932578, DOI: 10.1155/2013/932578.
- [14] H. Eltayeb and A. Kiliçman, A note on solutions of wave, Laplace's and heat equations with convolution terms by using a double Laplace transform, *Applied Mathematics Letters* **21**(12) (2008), 1324 – 1329, DOI: 10.1016/j.aml.2007.12.028.
- [15] H. Eltayeb and A. Kiliçman, On double Sumudu transform and double Laplace transform, *Malaysian Journal of Mathematical Sciences* **4**(1) (2010), 17 – 30, DOI: 10.1134/S1995080209030044.
- [16] A. P. Hiwarekar, Extension of prefunctions and its relation with Mittag-Leffler function, *The Journal of Analysis* **28** (2020), 169 – 177, DOI: 10.1007/s41478-017-0052-7.
- [17] A. P. Hiwarekar, New mathematical modeling for cryptography, *Journal of Information Assurance and Security* **9** (2014), 27 – 33, URL: <http://www.mirlabs.org/jias/secured/Volume9-Issue1/Paper4.pdf>.
- [18] A. P. Hiwarekar, Triple Laplace transforms and its properties, *Advances and Applications in Mathematical Sciences* **20**(11) (2021), 2843 – 2851.
- [19] T. Kim and D. S. Kim, Degenerate Laplace transform and degenerate gamma function, *Russian Journal of Mathematical Physics* **24** (2017), 241 – 248, DOI: 10.1134/S1061920817020091.
- [20] Y. J. Kim, B. M. Kim, L.-C. Jang and J. Kwon, A note on modified degenerate gamma and Laplace transformation, *Symmetry* **10**(10) (2018), 471, DOI: 10.3390/sym10100471.
- [21] N. Kokulan and C. H. Lai, A Laplace transform method for the image in-painting, in: *12th International Symposium on Distributed Computing and Applications to Business, Engineering & Science* (Kingston upon Thames, UK, 2013), pp. 243 – 246, DOI: 10.1109/DCABES.2013.51.
- [22] G. V. Nozhak and A. A. Paskar, Application of a modified discrete Laplace transform to finding processes in certain hyperbolic system with distributed parameters (Russian), *Mathematical Methods in Mechanics. Mat. Issled* **57** (1980), 83 – 90.
- [23] S.-L. Nyeo and R. R. Ansari, Space Bayesian learning for the Laplace transform inversion in dynamic light scattering, *Journal of Computational and Applied Mathematics* **235**(8) (2011), 2861 – 2872, DOI: 10.1016/j.cam.2010.12.008.

- [24] A. D. Poularikas and S. Seely, Laplace transform, Chapter 5, in *The Transforms and Applications Handbook*, 2nd edition, Boca Raton, CRC Press LLC, (2000), URL: <https://dsp-book.narod.ru/TAH/ch05.pdf>.
- [25] E. J. Riekstyn's, On certain possibilities of solution of a system of telegraph equations using the Laplace transform in the case of a composite conductor (Russian), *Latvijas Valsts Univ. Zinatn. Raksti* **8** (1956), 49 – 53.
- [26] M. Saif, F. Khan, K. S. Nisar and S. Araci, Modified Laplace transform and its properties, *Journal of Mathematics and Computer Science* **21**(2) (2020), 127 – 135, DOI: 10.22436/jmcs.021.02.04.
- [27] J. L. Schiff, *The Laplace Transform: Theory and Applications*, Springer, New York, xiv + 236 pages (1999), DOI: 10.1007/978-0-387-22757-3.
- [28] G.-Y. Tsaur and J. Wang, Close connections between the methods of Laplace transform, quantum canonical transform, and supersymmetry shape-invariant potentials in solving Schrodinger equations, *Chinese Journal of Physics* **53**(4) (2015), DOI: 10.6122/CJP.20150330.
- [29] S. Viaggiu, Axial and polar gravitational wave equations in a de Sitter expanding universe by Laplace transform, *Classical Quantum Gravity* **34** (2017), 035018, DOI: 10.1088/1361-6382/aa5570.

